



STUDIES IN THE ANTIMICROBIAL ACTIVITIES OF SOME NEWLY SYNTHESIZED TRIAZINES BY DISC DIFFUSION METHOD AND MIC METHODS

Dr. A. S. Shendge, D. T. Tayade* and P. R. Kale

Department of Chemistry, Government Vidarbha Institute of Science and Humanities,
Amravati 444604 India.

Article Received on
27 May 2017,

Revised on 16 June 2017,
Accepted on 5 July 2017

DOI: 10.20959/wjpps20178-9705

*Corresponding Author

D. T. Tayade

Department of Chemistry,
Government Vidarbha
Institute of Science and
Humanities, Amravati
444604 India.

ABSTRACT

2-Thiol-3-ethyl-4-ethylamino-6-[2-isobutoxy-5-(4-methyl-5-carboxy-1,3-thiazo-2-yl)]-phenyl-1,3,5-triazine (**A6**), 2-thiol-3-ethyl-4-tert-butylamino-6-[2-isobutoxy-5-(4-methyl-5-carboxy-1,3-thiazo-2-yl)]-phenyl-1,3,5-triazine (**A7**), 2-thiol-3-ethyl-4-p-chlorophenylamino-6-[2-isobutoxy-5-(4-methyl-5-carboxy-1,3-thiazo-2-yl)]-phenyl-1,3,5-triazine (**A8**), 2-thiol-3-ethyl-4-o-tolylamino-6-[2-isobutoxy-5-(4-methyl-5-carboxy-1,3-thiazo-2-yl)]-phenyl-1,3,5-triazine (**A9**), 2-thiol-3-ethyl-4-m-tolylamino-6-[2-isobutoxy-5-(4-methyl-5-carboxy-1,3-thiazo-2-yl)]-phenyl-1,3,5-triazine (**A10**), i.e. were synthesized and its antimicrobial activities were tested against *E.Coli*, *B.Subtilis*, *S.Typhi*, *S.Aureus*, *P.Vulgaris*, *A.Aerogenes* by Disc

diffusion method and MIC methods. These compounds showed good results.

KEYWORDS: *E.Coli*, *B.Subtilis*, *S.Typhi*, *S.Aureus*, *P.Vulgaris*, *A.Aerogenes*.

INTRODUCTION

The interactions between potent chemicals and living system contribute to understand the life processes and provide effective methods for the treatment, prevention and diagnosis of many diseases. A chemical substances used for this purpose are referred as 'drugs' and their actions on living system are known as 'drug effect' which can be preliminary tested by counting the antimicrobial activities of any molecule. 1,3,5-triazines has been reported as potent antagonist's interaction in vitro and in cell-based assays and also proved that they possess anti-consultant, anxiolytic, anti-tumor properties.^[1] It is effective against

cholecystokinin receptor (CCK), opiate receptor and platelet glycoprotein antagonists.^[2-3] The triazines are widely used as sedative, anti-depressive, anti-inflammatory and hypnotic agents.^[4-5] It is also used as dyes for acrylic fibers.^[6] Hence it was thought interesting to carry out the antimicrobial activities against various microbes to check the drug properties of newly synthesized 1,3,5-triazines by Disc diffusion method and MIC methods in this laboratory.^[7,8,9,10]

EXPERIMENTAL

The antibacterial activities of A-6 to A-10 compounds were tested to evaluate their efficiencies against pathogenic organisms. All the chemical and media were purchased from M/s. Hi-Media Pvt. Ltd., Mumbai, India. The organisms used were taken for studies *E.Coli*, *B.Subtilis*, *S.Typhi*, *S.Aureus*, *A.Aerogenes*, *P.Vulgaris*.

For the evaluation of in-vitro antimicrobial activity, the following three conditions must be fulfilled,

- i) First the substance to be evaluated must be brought in an intimate contact with the test organisms against which activity is to be estimated.
- ii) Secondly, favorable conditions must be provided to offer a maximum opportunity for optimum growth of the organisms in absence of antimicrobial agent, and
- iii) Thirdly, there should be a method for measuring antibacterial response obtained by antimicrobial agent.

Various methods had been proposed and adopted for the measurement of antibacterial activity, these are,

1. Agar streak dilution method.
2. Agar diffusion method.
3. Turbidometric method.
4. Serial dilution method.

In the present study, we used agar disc diffusion method to find out the activity of all synthesized compounds against the microbes. Then the minimum inhibitory concentrations were measured by serial dilution method for those compounds only which were found to be active.

A) Media Used**1. Nutrient Agar Medium - Composition**

Yeast extract	--	1.5 gms
Beef extract	--	1.5 gms
Peptone	--	5.0 gms
NaCl	--	5.0 gms
Agar powder	--	20 gms
Distilled water	--	1000ml
pH	--	7.4±0.2(at25°C)

2. Nutrient Broth Medium - Composition

Yeast extract	--	1.5 gms
Beef extract	--	1.5 gms
Peptone	--	5.0 gms
NaCl	--	5.0 gms
Distilled water	--	1000 ml
pH	--	7.4±0.2(at25°C)

Both the above cited media used were of bacteriostatic grade. Above media were found to be suitable for the growth of all four organisms used in the present work.

B) Slant preparation

Nutrient agar medium was dissolved in distilled water and then sterilized by auto claving. About 5 ml of molten media was transferred aseptically in previously sterilized test tubes. The test tubes were then plugged tightly and placed in a slanting position to cool and solidify.

C) Stock culture

Culture was grown on nutrient agar slants by incubating them for 24 hrs at 37°C.

D) Culture dilution (Sub-culturing)

One loopful of stock culture was added to 5 ml of nutrient broth medium for inoculation. The inoculated broth was incubated for 24 hrs at 37°C. For all experimental purposes 24 hrs fresh diluted culture of both the organisms were used.

E) Preparation of sample solution

An antibacterial activity is usually tested by making aqueous solution samples. However, compounds used in the present study are insoluble in water. Hence, to study antimicrobial activity their dilutions were prepared by using ethanol. Thus, ethanol was taken and tested as control.

To check the potency of compounds, the solutions were prepared with 50 µg/ml concentration. 1 ml of this solution was added to 5 ml of nutrient broth solution containing organism to be tested. Tubes with organism and medium with solvent were used as controls. These tubes were kept for incubation at 37°C for 24 hrs. Most of the compounds under study exhibited total inhibition of the test cultures within 24 hrs of incubation. The tube containing compounds showing inhibition (antimicrobial activity) was clear and the tube which was kept as control where no compound was added showed growth. Therefore, for all the antibacterial screenings, the concentrations of 50 µg/ml was used, which is in the range of the substance to be used as antibiotic.

F) Disc diffusion method

Every time fresh sterile nutrient agar medium was prepared. The proceedings were carried out aseptically. All the glassware and apparatus required were sterilized. In each sterile Petridish 15-20 ml of molten medium was added. Simultaneously 0.05-0.1 ml (approx. 2-3 drops) of 24 hrs fresh diluted culture of organism under study was added to each petriplate. The nutrient broth culture and nutrient agar media were mixed thoroughly by rotatory motion of agar plate on a plane surface. It was allowed to solidify at room temperature. Then sterilized Whatmann filter paper No. 1 discs (6 mm diameter) thoroughly moistened with the same concentration of each of the compound were placed on the surface of the plate. Disc moistened with ethanol was used as control. They were allowed to diffuse in the media and then the plates were incubated at 37°C for 24 hrs. The diameter of the zones of inhibition was observed.

The same procedure was followed for determining antifungal activity; only the potato dextrose plate was used.

3. Potato dextrose agar - Composition

Potato infusion form	--	200 gms
Dextrose	--	20 gms

Agar	--	15 gms
Distilled water	--	1000 ml
pH	--	5.6±0.2(at25°C)

The compounds, which showed antimicrobial activity, were further tested for their minimum inhibitory concentration by Serial Dilution Method.

G) Serial dilution method

To determine the MIC of various compounds the following procedure (Serial Dilution Method) was followed.

Nutrient broth was prepared by dissolving 13 gms of dehydrated medium in 1 liter of distilled water. The pH of the medium was adjusted to 7.4. 5ml of the medium was distributed in each tube. All the tubes were sterilized at 121°C for 20 minutes.

The appropriate amount of test compound was dissolved in the solvent ethanol gave final concentration of 1×10^{-2} M. Various amounts of the above stock solution was aseptically added to the various nutrient broth tubes (viz. 0.5, 1.0, 1.2, 1.4, 1.6, 1.8, .0 5.8, 6.0 ml). Fresh culture of the test bacterium was inoculated in each tube (0.2 ml culture). The inoculum size of the test bacterium was adjusted to give approx. 10^7 cFu. All the tubes were incubated at 37°C for 24 hrs. Uninoculated tube was kept as a control in which nutrient broth and 5 ml of the solvent was taken.

After 24 hours of incubation, all the tubes were observed for MIC against test bacterium. This was observed by the absence of visual turbidity in the tube receiving the highest dilution of the test compounds. To determine MIC of various test compounds against moulds (fungus) the following procedure was adopted. Potato dextrose broth was prepared as follows. 200 gm. of potato (Peeled) was added to 1 liter of distilled water. It was steamed for 20 min and volume adjusted to 1 liter. 20 gm of dextrose was added to this.

Appropriate amount of test compounds were dissolved in ethanol mixture to gave final concentration of 1×10^{-2} M. Various amounts of the above stock solution was added aseptically to the potato dextrose broth tubes (viz. 0.5, 1.0, 1.2, .6 ml). Fresh fungal culture was inoculated aseptically in each tube (0.2 ml of culture). All the tubes were incubated at 28°C for 96 hrs. After 48 hrs. of incubation all the tubes were observed for the MIC of test compounds.

RESULTS AND DISCUSSION

Total 05 synthesized compounds were studied for their antimicrobial activities. All the pathogens tested during analysis are human pathogens. The activities of compounds were tested against all the pathogens by disc diffusion method. It was found that all the compounds are active against *E.Coli*, *B.Subtilis*, *S.Typhi*, *S.Aureus*, while inactive against *A.Aerogenes*, *P.Vulgaris* as shown in Table No-1. MIC values were measured for the active compounds only, and given in Table No.-2

Activity against *E. coli*

E. Coli is a gram negative parasite living only in human or animal intestine. The clinical infections caused by *E. coli* were urinary tract infection, diarrhea, pathogenic infection and septicemia. Generally the patients of diarrhea were observed in February to July. The treatment of these patients who suffers diarrhea, normally medicinal practitioners make use of sulphonamides, cotrimoxazole, quinolones, ampicillin, cloxacillin, ciperacillin, carbenicillin, cephalosporin, gentamycine, chloramphenicol, tetracycline etc. antibiotics. These drugs directly affect on digestive system and ultimately circulatory system and finally on kidneys. Also the above drugs once used for the treatment of *E. coli* infection should not be used upto six months.

The antimicrobial activity of the synthesized compounds against *E. coli* is highly remarkable, all compounds are highly active.

Table 1: Activity of 1,4-substituted triazines.

Comp.	<i>E. Coli</i>	<i>B. Subtilis</i>	<i>S. Typhi</i>	<i>S. Aureus</i>	<i>A. Aerogenes</i>	<i>P. Vulgaris</i>
A6	Active	Active	Active	Active	Inactive	Inactive
A7	Active	Active	Active	Active	Inactive	Inactive
A8	Active	Active	Active	Active	Inactive	Inactive
A9	Active	Active	Active	Active	Inactive	Inactive
A10	Active	Active	Active	Active	Inactive	Inactive

Table 2: MIC values of active compounds in mgml⁻¹.

Comp.	<i>E. Coli</i>	<i>B. Subtilis</i>	<i>S. Typhi</i>	<i>S. Aureus</i>
A-6	780	856	853	1135
A-7	660	469	383	784
A-8	650	1185	1795	1662
A-9	550	479	473	540
A-10	862	781	1045	1345

MIC values

Inactive; 3500-1900

Weakly active; 1800-1500 1400-1000

Moderately active; 1400-1000

Highly active. < 1000

Activity against *E. coli*

From, Table No.-2 it can be easily seen that the compounds A-8 and A-9 showed highly activity in minimum concentration. Compounds A-8 and A-9 contain 1,3,5-triazino nucleus along with 1,4-thiazole nucleus respectively these nuclei may be responsible for the higher activity of these compounds. So these synthesized drugs can be used as the best alternative drugs for the treatment of diseases caused by *E. coli*.

2-Thiol-3-ethyl-4-ethylamino-6-[2-isobutoxy-5-(4-methyl-5-carboxy-1,3-thiazo-2-yl)]-phenyl-1,3,5-triazine (A6), 2-thiol-3-ethyl-4-tert-butylamino-6-[2-isobutoxy-5-(4-methyl -5-carboxy-1,3-thiazo-2-yl)]-phenyl-1,3,5-triazine (A7), 2-thiol-3-ethyl-4-p-chlorophenyl-amino-6-[2-isobutoxy-5-(4-methyl-5-carboxy-1,3-thiazo-2-yl)]-phenyl-1,3,5-triazine (A8), 2-thiol-3-ethyl-4-o-tolylamino-6-[2-isobutoxy-5-(4-methyl-5-carboxy-1,3-thiazo-2-yl)]-phenyl-1,3,5-triazine (A9), 2-thiol-3-ethyl-4-m-tolylamino-6-[2-isobutoxy-5-(4-methyl-5-carboxy-1,3-thiazo-2-yl)]-phenyl-1,3,5-triazine (A10), All compounds are highly active against *B. Subtilis* except A-8 which is moderately active.

A-6, A-7 and A-9 compounds are highly active, A-10 is moderately active while, A-8 is weakly active against *S. Typhi*.

A-7 and A-9 compounds are highly active, A-6 and A-10 is moderately active while, A-8 is weakly active against *S. Aureus*.

ACKNOWLEDGEMENTS

Authors are thankful to the UGC New Delhi for giving fellowship to author.

REFERENCES

1. Kalinski C., Umkehrer M., Ross G., Kolb J., Burdacka, C., Hiller W., *Tetrahedron Lett.*, 2006; 47: 3423.
2. Hulme C., Peng J., Tang S. Y., Burns C. J., Morize I., Labaudiniere R. J., *Org. Chem.*, 1998; 63: 8021.
3. Keating T. A., Armstrong R. W., *J. Org. Chem.*, 1996; 61: 8935.
4. Aversa M. C., Ferazzo A., Giannetto P., Kohnke F. H., *Synthesis*, 1986; 230.

5. Chimirri A., Grasso S., Ottana R., Romeo G., Zappala M., *J. Heterocyclic Chem.*, 1990; 27: 371.
6. Herbert J. A. L., Suschitzky H. *J. Chem. Soc., Perkin Trans.*, 1974; 1: 2657.
7. Bansal R.K., *J. Heterocyclic Chemistry*, 2012; 8: 12-24.
8. Fernandes P.S and Sonar T.M., *J.Ind.Chem.Soc.*, 1986; 53(4): 427.
9. Saleem F., *Eur. Pat.*, CHAPPL 87/1 APR 13, 3600009 (1987), *Chem Abstr.*, 1989; 110: 114893.
10. Hedge J.C., Sathesha Rai N. and Balkrishna K., *J.Chem.Sci.*, 2007; III 9(4): 299-302.