



GC-MS PROFILING OF BIOACTIVE COMPOUNDS FROM TALINUM PORTULACIFOLIUM (FORSSK.) ASCH. EX SCHWEINF. LEAF

Hemalatha K.* and Abirami P.

PG and Research Department of Botany, Vellalar College for Women (Autonomous), Erode
638 012, Tamilnadu. India.

Article Received on
23 May 2018,
Revised on 13 June 2018,
Accepted on 03 July 2018
DOI: 10.20959/wjpps20188-12057

*Corresponding Author

Hemalatha K.

PG and Research
Department of Botany,
Vellalar College for Women
(Autonomous), Erode 638
012, Tamilnadu. India.

ABSTRACT

Talinum portulacifolium (Forssk.) Asch. ex Schweinf. is the valuable medicinal plant from the family portulacaceae. It is used as the common leafy vegetable in many countries. The present study aims to the Gas chromatography- Mass spectrum analysis of Ethanolic extract of *Talinum portulacifolium* leaf. It is used as antidiabetic, aphrodisiac and to cure mouth ulcers. Leaves are used to treat eye disorders. Root is used in treatment of cough, pulmonary tuberculosis and gastritis. The present study showed 30 important phytoconstituents like Isobutyl- (5- Morpholin- 4- Yl- 1,2,3,4- Tetrahydro- 7- Thia- 6,9,10,11- Tetraazabenzoc[*C*]fluoren-8- Yl)- Amine, 6,7-Bis(trimethylsilyl)-4- methoxy- 1,3- diazobiphen and this GC-MS

analysis give ideas to isolate the chemical compounds effectively.

KEYWORDS: *Talinum portulacifolium*, portulacaceae, GC-MS, phytoconstituents, chromatogram, 6,7-Bis(trimethylsilyl)-4-methoxy-1,3-diazabiphenylene, Squalene, Neophytadiene, Misenine.

INTRODUCTION

Medicinal plants have a strong linkage with human health. Indian system of medicine like Ayurveda, Yunani and Sidha depend on medicinal plants for herbal drugs. Traditionally used medicinal plants produce a variety of compounds of known properties.^[1] World health organization estimates that up to 80% of the world's population relies mainly on herbal medicine for primary health care. Herbal medicine, which is also known as folk medicine, is known from every continent.^[2] Medicinal plants are an important element of indigenous

medical systems in all over the world. The ethno botany provides a rich resource for natural drug research and development.^[3]

Talinum portulacifolium (Forssk.) Asch. Ex Schweinf is one of the medicinal plant under the family Portulacaceae. This family is cosmopolitan and has 19 genera and it is distributed from Rajasthan to the peninsular region.^[4,5] It is used in the treatment of arthritis, backache, diarrhoea and sexually transmitted diseases like syphilis.^[6] It is also used as a medicine for constipation and ulcer.^[7] The green leafy vegetables rich in antioxidants and contain varying amount of phytochemicals like vitamin C, flavonoids and carotenoids. The leafy vegetables are having rich chlorophyll content and it has been proved to build red blood cells and help to decrease the risk of heart disease, stroke, and certain cancers.^[8]

The knowledge of chemical constituents of plants would further be valuable in discovering the actual value of folkloric remedies.^[9] Gas Chromatography Mass Spectroscopy, a hyphenated system which is a very compatible technique and the most commonly used technique for the identification and quantification purpose. The unknown organic compounds in a complex mixture can be determined by interpretation and also by matching the spectra with reference spectra.^[10] The objective of the present study deals with the isolation of the chemical compounds from the ethanolic extract of *Talinum portulacifolium* leaf.

MATERIALS AND METHODS

Collection and identification

The fresh plant material was collected from Pungambadi, Erode District, Tamilnadu, India. The plant material was identified by using the local floras.^[11,12] The plant was confirmed with the help of type specimens available in the Herbarium of Botanical Survey of India, Southern region center, Coimbatore, Tamil Nadu. The Herbarium number in BSI is **BSI/SRC/5/23/2016/Tech./1358**. The herbarium was deposited in the Vellalar College for Women (Autonomous) Herbarium, Erode 638 012, Tamilnadu.

Gas Chromatography-Mass Spectrometry Analysis

GC-MS analysis of Ethanolic extract of the leaf of *Talinum portulacifolium* (Forssk.) Asch. ex Schweinf. was performed using a GC-MS (Model; Thermo Trace GC Ultra Ver.5.0) equipped with a DB-35MS fused silica capillary column (30 m length X Outside diameter 0.25 mm X internal diameter 0.25 µm) and gas chromatogram interfaced to a Mass selective detector (MS-DSQ- II) with XCALIBUR software. For GC-MS detection, an electron

ionization system with ionization energy of -70eV was used. Helium gas was used as a carrier gas at a constant flow rate of 1ml/min and the sample injected was $1\mu\text{l}$; Injector temperature was 250°C ; Ion source temperature was 200°C . The oven temperature was programmed from 70° to 200°C at the rate of 10°C/min , held isothermal for 1 minutes and finally raised to 250°C at 10°C/min . Interface temperature was kept at 250°C . The relative percentage of each extract constituent was expressed as percentage with peak area normalization.

Identification of compounds

The identity of the compounds in the extract was assigned by the comparison of their retention time and mass spectra fragmentation patterns with those stored on the computer library and also with published literatures. NIST^[13], WILEY^[14] library sources were also used for matching the identified compounds from the plant material.

RESULTS

The GC-MS analysis of ethanolic extract of *Talinum portulacifolium* (Forssk.) Asch. ex Schweinf. revealed the presence of thirty compounds. The GC-MS running time was 40.51 minutes. The GC-MS chromatogram of *Talinum portulacifolium* is presented in Fig.1. The active principles with their Retention Time (RT), Molecular formula, Molecular weight (MW) and Peak area are presented in the Table.1. The spectra of the compounds are matched with Wiley 9.0 and NIST libraries.

The most prevailing major compounds of *Talinum portulacifolium* are Isobutyl-(5-morpholin-4-yl-1,2,3,4-tetrahydro-7-thia-6,9,10,11-tetraaza-benzo[c]fluoren-8-yl)-amine (13.11%), 6,7-Bis(trimethylsilyl)-4-methoxy-1,3-diazabiphenylene (11.64%), 2,8-Dimethyl-2-(4,8,12-trimethyltridecyl)-6-chromanol (11.41%), Methyltris(trimethylsiloxy)silane (8.95%), Phytol (6.64%), Cyclohexane, 1,1',1'',1'''-(1,6-hexanediylidene)tetrakis(CAS) (5.18%), Neophytadiene (5.14%), Squalene (4.23%), Misenine (4.18%), 4,8,12,16-Tetramethylheptadecan-4-olide (0.41%), (E,E)-Farnesylacetone (0.43%), 11,13-Tetradecadien-1-ol acetate (0.48%), Dinaphtho[2,1-b:1',2'-d]thiophene (CAS) (0.50%), Lucenin 2 (0.58%), Silane, triethoxymethyl- (CAS) (0.62%) are also detected in minor amount.












Figure 1: Chromatogram of Ethanolic Extract of *Talinum portulacifolium* (Forssk.) Asch. ex Schweinf.

Table 1: Phytochemicals Identified In the Ethanolic Extract of *Talinum portulacifolium* (Forssk.) Asch. ex Schweinf. LEAF.

S.NO.	R.T	COMPOUND NAME	MOLECULAR FORMULA	MOLECULAR WEIGHT	AREA %
1	3.19	Silane, triethoxymethyl- (CAS)	C7H18O3Si	178	0.62
2	4.47	4-Isopropoxy-5-methoxybenzo[b]furan	C12H14O3	206	0.89
3	4.78	Silane, diethoxydimethoxy-	C6H16O4Si	180	1.31
4	7.73	2-Methylenecyclohexan-1-carboxylic acid	C14H15ClO2	250	2.02
5	10.70	Tetradecene	C14H28	196	2.23
6	13.08	2-tert-Butyl-4-isopropyl-5-methylphenol	C14H22O	206	3.61
7	14.85	Cetene	C16H32	224	1.80
8	19.13	5-Octadecene, (E)-	C18H36	252	2.20
9	20.00	Neophytadiene	C20H38	278	5.14
10	20.50	3,7,11,15-Tetramethyl-2-hexadecen-1-ol	C20H40O	296	1.02
11	20.86	3,7,11,15-Tetramethyl-2-hexadecen-1-ol	C20H40O	296	1.70
12	21.54	11,13-Tetradecadien-1-ol acetate	C16H28O2	252	0.48
13	22.39	Methyl 1,3-dihydro-2H-isobenzofuran-4-carboxylate	C10H10O3	178	0.56
14	23.14	Hexadecanoic acid, ethyl ester (CAS)	C18H36O2	284	1.03
15	25.31	Phytol	C20H40O	296	6.64
16	29.29	4,8,12,16-Tetramethylheptadecan-4-olide	C21H40O2	324	0.41
17	29.59	(E,E)-Farnesylacetone	C18H30O	262	0.43
18	30.56	Dinaphtho[2,1-b:1',2'-d]thiophene (CAS)	C20H12S	284	0.50
19	31.04	1-Dodecanol (CAS)	C12H26O	186	0.68
20	31.48	Lucenin 2	C27H30O16	610	0.58
21	32.42	Methyltris(trimethylsiloxy)silane	C10H30O3Si4	310	8.95
22	32.99	2,3,9,10-Tetramethoxy-6,13-dithia-5,7,12,14-tetraazap	C20H16N4O4S2	440	3.23
23	33.54	(5E)-2,2,7-Trimethyl-4-oxaocta-5,7-dienal	C10H16O2	168	3.19

24	34.85	Isobutyl-(5-morpholin-4-yl-1,2,3,4-tetrahydro-7-thia-6,9,10,11-tetraaza-benzo[c]fluoren-8-yl)-amine	C ₂₀ H ₂₆ N ₆ O ₅	398	13.11
25	35.68	Cyclohexane, 1,1',1'',1'''-(1,6-hexanediyldiene)tetrakis(CAS)	C ₃₀ H ₅₄	414	5.18
26	36.21	Squalene	C ₃₀ H ₅₀	410	4.23
27	37.41	(22R,24S)-24,25-epoxy-22-acetoxy-7 α -hydroxyergosta-1,4-dien-3-one	C ₃₀ H ₄₄ O ₅	484	1.03
28	38.47	Misenine	C ₃₃ H ₅₄ N ₂ O	494	4.18
29	39.00	6,7-Bis(trimethylsilyl)-4-methoxy-1,3-diazabiphenylene	C ₁₇ H ₂₄ N ₂ O ₂ Si ₂	328	11.64
30	39.51	2,8-Dimethyl-2-(4,8,12-trimethyltridecyl)-6-chromanol	C ₂₇ H ₄₆ O ₂	402	11.41

Table 2: Structure and Hit Spectrum of some important Biocompounds.

S.No	Compound name	Structure	Hit Spectrum
1.	Isobutyl-(5-Morpholin-4-Yl-1,2,3,4-Tetrahydro-7-Thia-6,9,10,11-Tetraaza-Benzo[C]Fluoren-8-Yl)-Amine		
2.	6,7-Bis(trimethylsilyl)-4-methoxy-1,3-diazabiphenylene		
3.	2,8-Dimethyl-2-(4,8,12-Trimethyltridecyl)-6-Chromanol		
4.	Methyltris(trimethylsiloxy) Silane		

The highest peak area percentage of 13.11% was obtained by Isobutyl-(5-morpholin-4-yl-1,2,3,4-tetrahydro-7-thia-6,9,10,11-tetraaza-benzo[c]fluoren-8-yl)-amine. Lowest peak area

percentage was obtained by 4,8,12,16-Tetramethylheptadecan-4-olide (0.41%). Some important biocompounds structure and hit spectrum are displayed in the Table.2.

DISCUSSION

The important phytoconstituent phytol present in the present study was also reported in *Talinum paniculatum* (jacq.) Gertn.^[15] The GC-MS analysis of *Talinum portulacifolium* was studied and the major chemical constituents from the column fraction were Dodecane, Tetradecamethyl cycloheptasiloxane, Hexadecanoic acid (methyl ester), cyclopentadiene.^[16]

The important biocompounds were reported in the GC-MS analysis of the whole plant *Hydrocotyle javanica* like 3,7,11,15- Tetramethyl-2-hexadecen-1-ol, Phytol, Squalene.^[17] These same compounds were identified in the present study. Important biocompound Octadecane was identified in the GC-MS analysis of *Portulaca oleracea*. The same compound was identified in the present analysis.^[18]

CONCLUSION

The present study showed the GC-MS analysis of the ethanolic extract of *Talinum portulacifolium* leaf. There were thirty compounds present in the extract. These compounds are the reason for the biological activity of the plant. These bioactive compounds may be responsible for their efficacy and the healing property of various diseases.

ACKNOWLEDGEMENT

We are thankful to the Medical Textiles – Analytical Division, The South India Textile Research Association, Coimbatore 641014, Tamil Nadu, India.

Conflict of Interest

Declared none.

REFERENCES

1. Haeborne SB, and Baxter H, Phytochemical dictionary. A hand book of bioactive compounds from plants. Taylor and Francis, London, 1995.
2. BGCI Fact sheet.2000. Plants as medicine. Botanic garden conservation International/ British Airways/ International centre for conservation education. www.wwf.org.uk/filelibrary/pdf/useofplants.pdf.
3. Farnsworth NR. The role of ethno pharmacology in drug development. Ciba Found Symp., 1990; 154: 2-11.

4. Heywood VH, U.K: Oxford University Press. Flowering plants of the World, 1978; 336.
5. Anonymous, The wealth of India- A dictionary of Indian raw materials and industrial products. 1974. Raw materials Vol.10. CSIR, New Delhi, p.113.
6. Gerber M, Boutron-Ruault MC, Hercberg S, Riboli E, Scalbert A and Siess MH, Food & cancer: state of the art about the protective effect of fruits and vegetables. *Bulletin du Cancer.*, 2002; 89(3): 293-312.
7. John Warui Kiring, A survey of Traditional Health remedies used by the Maasai of Southern Kaijiado District, Kenya, *Ethanobotany Research and Applications*, 2006; 4: 061-073.
8. Mojib F, Kamalinejad M, Ghaderi N and Vahidipour HR. Phytochemical screening of some species of Iranian plants. *Iranian journal of Pharma Research*, 2003; 2(2): 77-82.
9. Thangavel K, Maridass M, Sasikala and M, Ganesan V. *In vitro* Micropropagation of *Talinum portulacifolium* L. through Axillary Bud Culture, *Ethnobotanical Leaflets*, 2008; Pp. 413-418.
10. Ronald Hites A. Gas Chromatography Mass Spectroscopy: Handbook of Instrumental Techniques for Analytical Chemistry, 1997; 609-611.
11. Gamble JS. and Fisher, C.E.C. Flora of the Presidency of Madras. London: Ad. Lord and Sons Limited, 1956; I-III.
12. Mathew KM. The Flora of Tamilnadu Carnatic. Tamilnadu. 1983, The herbarium, St. Joseph's College.
13. Mc Lafferly, F.W. Registry of mass spectral data, 5 ed, Wiley New York: John Wiley & Sons Inc., 1989.
14. Stein, SE, National Institute of Standards and Technology (NIST) Mass Spectral Data base, 1990. Version 3.02, Gaithersburg Md. USA.
15. Thanamool C, Papirom P, Chanlun S, and Kupittayanant S. *Talinum paniculatum* (Jacq.) Gertn: A Medicinal Plant with potential Esterogenic activity in overiectomized Rats. *Inter. J. Pharma and Pharmaceu Sci.*, 2013; 5(2): Pp478- 485.
16. Vani M, Rahaman SKA and Rani AP. *In Vitro, In vivo* Antiasthmatic studies of *Talinum portulacifolium* F. *Asian. J. of Pharma.*, 2017; 11(2): S289-S301.
17. Namasivayam K, Karuppanan A, Habibulla AK., and Rangasamy N. 2017. GC- MS Profiling of Ethnomedicinal Plant *Hydrocotyle javanica* Thunb. *Oriental J Chem.*, 33(4): 2127-2133.

18. Hifnawy S.M., Mohammed R. and Montaser M. 2016. Chemical composition and antimicrobial activity of the lipoid extract from *Portulaca oleracea* (L.) plant, Family Portulacaceae. *World J. of pharma and pharma sci.*, 5(6): 2101-2108.