



EVALUATION OF IN VITRO ANTIUROLITHIATIC ACTIVITY OF *MAGNOLIA CHAMPACA*

Sai Sruthi G.* and Krishna Priyanka B.

Department of Pharmacognosy, Vishnu Institute of Pharmaceutical Education and Research,
Narsapur, Medak, Telangana, India.

Article Received on
01 Jan. 2019,

Revised on 21 Jan. 2019,
Accepted on 12 Feb. 2019

DOI: 10.20959/wjpps20193-12844

*Corresponding Author

Sai Sruthi G.

Department of
Pharmacognosy, Vishnu
Institute of Pharmaceutical
Education and Research,
Narsapur, Medak,
Telangana, India.

ABSTRACT

The present study was undertaken to evaluate the in vitro antiurolithiatic activity of the medicinal plant *Magnolia champaca*. Ethanolic extract showed their maximum efficiencies in the dissolution of calcium oxalate crystals. Our results have clearly indicated that the Ethanolic leaf extracts of *Magnolia champaca* were quite promising for further studies in this regard. In this study Neeri was used as standard drug.

KEYWORDS: In Vitro Antiurolithiatic Activity, Ethanolic Extract, Urolithiasis, *Magnolia Champaca*, Neeri.

1. INTRODUCTION

Urolithiasis is the third most common disorder of the urinary tract, is defined as the formation of sediment in the urinary tract consisting of one or more of the poorly soluble crystalloids of urine. It is a worldwide problem particularly common in parts of United States, South Africa, India and South East Asia. Approximately 2% of the world population experiences renal stone disease with a male-female ratio of 2:1 and the peak incidence is observed in 2nd to 3rd decade of life. Renal calculi are characterised clinically by colicky pain (renal colic) as manifest by hematuria. Major risk factors responsible for the nephrolithiasis are inadequate urinary drainage, microbial infections, diet with excess oxalates and calcium, vitamin abnormalities i.e.; deficiency of Vitamin-A, excess of vitamin D, metabolic diseases like hyperparathyroidism, cystinuria, gout, intestinal dysfunction.^[1]

Urolithiasis refers to the solid nonmetallic minerals in the urinary tract. Among the several types of kidney stones, the most common are calcium oxalate. Urolithiasis is a complex

process that is a consequence of an imbalance between promoters and inhibitors in the kidneys.^[2]

Nephrolithiasis or renal stone disease remains a significant health problem in the adult population, with serious medical consequences, throughout a patient's lifetime. The worldwide incidence of urolithiasis is quite high, and more than 80% of urinary calculi are calcium oxalate stones alone or calcium oxalate mixed with calcium phosphate.^[3]

The size and nature of crystals governs overall clinical manifestations of this complaint whereas urinary chemistry is one of the important factors in determining the type of crystals formed and the nature of macromolecules included on the surface of the crystals. Calcium oxalate stones make up the majority as they account for 70-80% of all kidney stones. Surgical operation, lithotripsy and local calculus disruption using high-power laser are widely used to remove the calculi. However, these procedures are highly costly and with these procedures recurrence is quite common.^[4] Various factors such as age, sex, industrialization, socioeconomic status, diet and environment, influences urolithiasis in terms of both site, and the physico-chemical composition of the calculi.^[5]

Thus a drug for the prevention of this disease or its recurrence would be of great interest. Medicinal plants have played a significant role in various ancient traditional systems of medication. Even today, plants provide a cheap source of drugs for majority of world's population. Several pharmacological investigations on the medicinal plants used in traditional antiurolithic therapy have revealed their therapeutic potential in the in vitro models ~ 19 ~ Journal of Medicinal Plants Studies.^[6-7]

Calcium containing stones may be in the form of pure calcium oxalate(50%) or calcium phosphate(5%) and a mixture of both(45%) followed by magnesium phosphate(15-20%),uric 5 acid(10%) and cystine(1%).^[8]

Magnolia champaca, known in English as champak is a large evergreen tree in the Magnoliaceae family. It was previously classified as *Michelia champaca*. It is known for its fragrant flowers, and its timber used in woodworking.^[9,10]

"*Magnolia champaca* however is more rare and has a strong perfume, and is not that commonly or plentifully used - for example in hair it is worn singly or as a small corsage but rarely as a whole garland, and for bridal beds it is most often jasmine and roses while for

bowls of water to be placed around rooms usually other, more colourful for visual decoration and less strongly perfumed flowers are used." [11] The scent similar to the scent of this plant is said to emit by a civet in Sri Lanka, *Paradoxurus montanus*. Because all the other civets are known to emit very unpleasant odours, this species is renowned to emit pleasant odour similar to this plant's scent. [12]

2. MATERIALS AND METHODS

Plant Material

The leaves of *Magnolia champaca* was collected in the month of October from Narsapur village, Medak dist. of Telangana, India. The plant was authenticated by D.Venkateshwara Rao, Deputy Director, Telangana. Forest Academy, Dullapally, Hyderabad, Rangareddy District. The leaves were washed with tap water and dried under shade.

Preparation of Plant Extract

The leaves were shade dried and powdered. The crude plant extract was prepared by Soxhlet extraction method. 50g of powdered plant material was extracted with 500ml of ethanol and water individually. The process of extraction was carried out up to 6 cycles, till the solvent in siphon tube of an extractor became colorless. The two extracts were filtered separately, and evaporated to dryness using rotary evaporator. Further the dried extracts were maintained in a refrigerator at 4°C for further antiurolithiatic activity.

Chemicals Used: Neeri, Sodium oxalate, Tris buffer, calcium chloride, Potassium permanganate (KMnO₄), Sulphuric acid (H₂SO₄).

Investigation of in Vitro Antiurolithiatic Activity Test By Titrimetry

The experimental kidney stones of calcium oxalate (CaOx) were prepared in the laboratory by taking equimolar solution of calcium chloride dehydrate in distilled water and sodium oxalate in 10 ml of 2N H₂SO₄. Both were allowed to react in sufficient quantity of distilled water in a beaker, the resulting precipitate was calcium oxalate. The precipitate was freed from traces of sulphuric acid by ammonia solution, washed with distilled water and dried at 60°C. The dissolution percentage of calcium oxalate was evaluated by taking exactly 1 mg of calcium oxalate and 10 mg of the extract, packed it together in semi permeable membrane of egg as shown in the model designed given below. This was allowed to suspend in a conical flask containing 100 ml of 0.1M Tris buffer. First group served as blank containing only 1 mg of calcium oxalate. The second group served as positive control containing 1 mg of calcium

oxalate and along with the 10mg standard drugs, i.e. Neeri. The 3rd group along with 1 mg of calcium oxalate contain Ethanolic extract. The conical flasks of all groups were kept in an incubator preheated to 37°C for 2 h. Remove the contents of semi permeable membranes from each group into separate test tubes, add 2 ml of 1N sulphuric acid to each test tube and titrated with 0.9494 N KMnO₄ till a light pink colour end point obtained. The amount of remaining undissolved calcium oxalate is subtracted from the total quantity used in the experiment in the beginning to know the total quantity of dissolved calcium oxalate by various solvent extracts.^[13]

3. RESULTS AND DISCUSSION

Drug therapy has developed in response to population health care^[14] needs. There are many crucial areas in medicine such as liver diseases, arthritis, old age related problems, certain viral infections and cancer where the conventional medicine is devoid of satisfactory treatment. These are among the promising areas of research and development of medicines from the vast highly potential plant resources. Plants are also attractive sources for the development of novel and very effective and safe therapeutic agents against kidney procumbens. Herbal medicines are also in great demand in the developed world for primary health care because of their efficacy, safety and lesser side effects¹⁵. Unlike allopathic medicines which target is only one aspect of urolithiatic pathophysiology, most of plant based therapy have been shown to be effective at different stages of stone pathophysiology¹⁶. About 80% of the world populations rely on the use of traditional medicine which is predominantly based on plant materials.^[17] Plant based drug discovery programmes continue to provide an important source of new drug leads.^[18] Lithiasis (stone formation) is an important cause for acute and chronic renal failure, includes both nephrolithiasis (stone formation in kidney) and urolithiasis (stone formation in ureter or bladder or both). Among the various kinds of stones identified, calcium stones occur mainly in Men, while phosphate stones formation is more in women.^[19]

This study evaluates the antiurolithiatic activity of Ethanolic leaf extract of *Magnolia champaca*. The highest percentage i.e. 97.1% of calcium oxalate {CaOx} dissolution was observed in Ethanolic extract. Ethanolic extract of *Magnolia champaca* was found to be more effective in dissolution of calcium oxalate than standard drug Neeri. From this study, it was observed that Ethanolic leaf extract of *Magnolia champaca* showed their highest dissolution of calcium oxalate. This study has given primary evidence for *Magnolia champaca* as the

plant which possess lithotriptic property. This in vitro study has given lead data and shown that Ethanolic leaf extract of *Magnolia champaca* was quite promising for further studies in this regard.

Table 1: Shows % dissolution of calcium oxalate (CaOx) by *Magnolia champaca* leaves extracts.

S.No	% of dissolution of calcium oxalate	
	Groups	<i>Magnolia champaca</i>
1.	Blank	0
2.	Positive Control	81
3.	Ethanolic extract	97.1



Figure 1(a): Decalcification of egg shell in 10% Acetic acid overnight.



Figure 1(b): Decalcified Eggs




Figure 1(c): Egg membrane along with the contents suspended into the 0.1 M Tris buffer.

Figure 1: In vitro experimental model setup to evaluate antiurolithiatic activity.

4. CONCLUSION

In vitro urolithiasis has been performed on the selected plant *Magnolia champaca* by using the standard drug, Neeri. The work was performed by using in vitro antiurolithiatic model for calculating percentage dissolution of kidney stone. Ethanolic leaf extract of *Magnolia champaca* shows highest dissolution than standard drug Neeri. This study has given primary evidence for *Magnolia champaca* as the plant which possess antiurolithiatic property.

ACKNOWLEDGEMENT

We sincerely thankful to our principal Dr. A.Ramesh and staff members, Director and chairman of our college Vishnu Institute of Pharmaceutical Education and Research (VIPER) for supporting us.

REFERENCES

1. HG Tiselius. Aetiology and investigation of stone disease. Curriculum in urology. Eur J Urol. 1998; 2(1): 1–7.
2. Khan SR. Interactions between stone-forming calcific crystals and macromolecules. Urol Int. 1997; 59: 59–71.
3. Mitra SK, Gopumodhavan S, Venkataranganna MV, Sundaran R. Effect of cystone, a herbal formulation on glycolic acid-induced urolithiasis in rats. Phytother Res., 1998; 12: 372–4.
4. Prasad K, Sujatha D and Bharathi K. Herbal drugs in urolithiasis – a review. Phcog Rev., 2007; 1: 175–179.

5. Moe OW. Kidney stones: Pathophysiology and medical management. *Lancet*, 2006; 367: 333-444.
6. Baumann JM. Stone prevention: why so little progress? *Urological Research*, 1998; 26: 77-81.
7. Leonetti F, Lechevallier E, Dussol B, Berland Y. Lithiase Urinaire—La Revue du Praticien, 1996; 26: 1557-1567.
8. Radha singanallur Ramu, Ravi Doraiswamy and Hiran Mai Yadav (2017). Antiurolithiatic activity of Aqueous bark extract of *Crateva Magna Lour.* (DC). *International Journal of Research in Ayurveda and Pharmacy*, 2017; 8: 271-278.
9. efloras.org: Flora of China treatment of *Michelia (Magnolia) champaca*. accessed 7.12.2015
10. Pacific Horticulture Society: "Striving for Diversity: Fragrant Champaca". accessed 7.12.2015
11. Kokate, C.K., Purohit, A.P. and Gokhale S.B. (2003), *Pharmacognosys*, 2nd edition, published by Nirali prakashan, pune, 2003; 522-523.
12. <http://wizzley.com/sri-lankan-brown-palm-civets-paradoxurus-montanus-non-ringtails-of-sri-lanka/>.
13. Minter, S. "Fragrant Plants." in Prance, G. and M. Nesbitt. (2005). *The Cultural History of Plants*. London: Routledge. 242. This is great.
14. Michael Dickson and Jean paul Gagnon. Key factors in the rising cost of new drug discovery and development. *Nature Reviews Drug Discovery*, 2004; 3: 417-429.
15. Kamboj V. P. Herbal medicine. *Current Science Association (JSTOR)*, 2000; 78(1): 35- 39.
16. Archana R. Dhole, Vikas R. Dhole, Chandrakant S. Magdum, Shreenivas Mohite. Herbal Therapy for Urolithiasis: A Brief Review. *Research Journal of Pharmacology and Pharmacodynamics*, 2013; 5(1): 6-11.
17. Subramoniam. A, P. Pushpangadan. Development of phytomedicines for liver disease. *Indian Journal of Pharmacology*, 1999; 31(3): 166-175.
18. Sanjay M, Jachak and Arvind saklani. Challenges and opportunities in drug discovery from plants. *Current Science Association (JSTOR)*, 2007; 92(9): 1251-1257.
19. Ramachandran. S, Vijayakumar. T. M, Sai sandeep. V, Ramsai. K and Dhanaraju. M. D. Antilithiatic Activity of Poly Herbal Extracts on Ethylene glycol-Induced Lithiasis in Rats. *European Journal of Biological Sciences*, 2011; 3(2): 36-39.